

2°. There exists a principal filter  $\mathcal{F}$  such that  $S = \partial\mathcal{F}$ .

$$\begin{aligned} \bigsqcup^{\mathfrak{A}} T \in \uparrow\uparrow S &\Leftrightarrow \text{up} \bigsqcup^{\mathfrak{A}} T \in S \Leftrightarrow \forall K \in \text{up} \bigsqcup^{\mathfrak{A}} T : K \in \partial\mathcal{F} \Leftrightarrow \\ &\forall K \in \text{up} \bigsqcup^{\mathfrak{A}} T : K \not\prec \mathcal{F} \Leftrightarrow \bigsqcup^{\mathfrak{A}} T \not\prec \mathcal{F} \Leftrightarrow \bigsqcup^{\mathfrak{A}} T \in \star\mathcal{F} \Leftrightarrow \exists \mathcal{K} \in T : \mathcal{K} \in \star\mathcal{F} \Leftrightarrow \\ \exists \mathcal{K} \in T : \mathcal{K} \not\prec \mathcal{F} &\Leftrightarrow \exists \mathcal{K} \in T \forall K \in \text{up} \mathcal{K} : K \not\prec \mathcal{F} \Leftrightarrow \exists \mathcal{K} \in T \forall K \in \text{up} \mathcal{K} : K \in \partial\mathcal{F} \Leftrightarrow \\ &\exists \mathcal{K} \in T : \text{up} \mathcal{K} \subseteq S \Leftrightarrow \exists \mathcal{K} \in T : \mathcal{K} \in \uparrow\uparrow S \Leftrightarrow T \cap \uparrow\uparrow S \neq \emptyset. \\ \perp \in \uparrow\uparrow S &\Leftrightarrow \text{up} \perp \subseteq S \Leftrightarrow \perp \in S \text{ what is false.} \end{aligned}$$

□

COROLLARY 1480. If  $S$  is a complete free star on  $\mathfrak{F}$  then  $\Downarrow S$  is a complete free star on  $\mathfrak{B}$ , provided that  $\mathfrak{B}$  is a complete lattice.

### 18.3.3. Complete staroids and multifunctors.

DEFINITION 1481. Consider an indexed family  $\mathfrak{B}$  of posets. A pre-staroid  $f$  of the form  $\mathfrak{B}$  is *complete* in argument  $k \in \text{arity } f$  when  $(\text{val } f)_k L$  is a complete free star for every  $L \in \prod_{i \in (\text{arity } f) \setminus \{k\}} \mathfrak{B}_i$ .

DEFINITION 1482. Consider an indexed family  $(\mathfrak{A}_i; \mathfrak{B}_i)$  of filtrators and pre-multifunctor  $f$  is of the form  $\prod \mathfrak{B}$ . Then  $f$  is *complete* in argument  $k \in \text{arity } f$  iff  $\langle f \rangle_k L \in \mathfrak{B}_k$  for every family  $L \in \prod_{i \in (\text{arity } f) \setminus \{k\}} \mathfrak{B}_i$ .

PROPOSITION 1483. Consider an indexed family  $(\mathfrak{F}_i; \mathfrak{B}_i)$  of primary filtrators over boolean lattices. Let  $f$  be a pre-multifunctor of the form  $\mathfrak{F}$  and  $k \in \text{arity } f$ . The following are equivalent:

- 1°. Pre-multifunctor  $f$  is complete in argument  $k$ .
- 2°. Pre-staroid  $\Downarrow [f]$  is complete in argument  $k$ .

PROOF. Let  $L \in \prod \mathfrak{B}$ . We have  $L \in \text{GR } [f] \Leftrightarrow L_i \not\prec \langle f \rangle_i L|_{(\text{dom } L) \setminus \{i\}}$ ;  
 $(\text{val } [f])_k L = \partial \langle f \rangle_k L$  by the theorem 1279.

So  $(\text{val } [f])_k L$  is a complete free star iff  $\langle f \rangle_k L \in \mathfrak{B}_k$  (proposition 1474) for every  $L \in \prod_{i \in (\text{arity } f) \setminus \{k\}} \mathfrak{B}_i$ . □

EXAMPLE 1484. Consider functor  $f = \text{id}^{\text{FCD}(U)}$ . It is obviously complete in each its two arguments. Then  $[f]$  is not complete in each of its two arguments because  $(\mathcal{X}; \mathcal{Y}) \in [f] \Leftrightarrow \mathcal{X} \not\prec \mathcal{Y}$  what does not generate a complete free star if one of the arguments (say  $\mathcal{X}$ ) is a fixed nonprincipal filter.

THEOREM 1485. Consider a semifiltered, star-separable, down-aligned filtrator  $(\mathfrak{A}; \mathfrak{B})$  with finitely meet closed and separable core where  $\mathfrak{B}$  is a complete boolean lattice and both  $\mathfrak{B}$  and  $\mathfrak{A}$  are atomistic lattices.

Let  $f$  be a multifunctor of the aforementioned form. Let  $k, l \in \text{arity } f$  and  $k \neq l$ . The following are equivalent:

- 1°.  $f$  is complete in the argument  $k$ .
- 2°.  $\langle f \rangle_l (L \cup \{(k; \bigsqcup X)\}) = \bigsqcup_{x \in X} \langle f \rangle_l (L \cup \{(k; x)\})$  for every  $X \in \mathcal{P} \mathfrak{B}_k$ ,  $L \in \prod_{i \in (\text{arity } f) \setminus \{k, l\}} \mathfrak{B}_i$ .
- 3°.  $\langle f \rangle_l (L \cup \{(k; \bigsqcup X)\}) = \bigsqcup_{x \in X} \langle f \rangle_l (L \cup \{(k; x)\})$  for every  $X \in \mathcal{P} \mathfrak{A}_k$ ,  $L \in \prod_{i \in (\text{arity } f) \setminus \{k, l\}} \mathfrak{B}_i$ .

PROOF.

3°  $\Rightarrow$  2°. Obvious.